

AGING ON PARISI'S TREE

J-P. Bouchaud and D.S. Dean

Service de Physique de l'Etat Condensé

Direction des Recherches sur l'Etat Condensé, les Atomes et les Molecules

Commissariat à l'Energie Atomique, Orme des Merisiers

91191 Gif-sur-Yvette CEDEX, France

(February 1, 2008)

Abstract

We present a detailed study of simple ‘tree’ models for off equilibrium dynamics and aging in glassy systems. The simplest tree describes the landscape of a random energy model, whereas multifurcating trees occur in the solution of the Sherrington-Kirkpatrick model. An important ingredient taken from these models is the exponential distribution of deep free-energies, which translate into a power-law distribution of the residence time within metastable ‘valleys’. These power law distributions have infinite mean in the spin-glass phase and this leads to the aging phenomenon. To each level of the tree are associated an overlap and the exponent of the time distribution. We solve these models for a finite (but arbitrary) number of levels and show that a two level tree accounts very well for many experimental observations (thermoremanent magnetisation, a.c susceptibility, second noise spectrum....).

We introduce the idea that the deepest levels of the tree correspond to equilibrium dynamics whereas the upper levels correspond to aging. Temperature cycling experiments suggest that the borderline between the two is temperature dependent. The spin-glass transition corresponds to the temperature at which the uppermost level is put out of equilibrium but is subsequently

followed by a sequence of (dynamical) phase transitions corresponding to non equilibrium dynamics within deeper and deeper levels. We tentatively try to relate this ‘tree’ picture to the real space ‘droplet’ model, and speculate on how the final description of spin-glasses might look like.

I. INTRODUCTION

Aging experiments in glasses and spin-glasses [1], [2], [3] are now the focus of an intense theoretical [4], [5], [6], [7], [8] and numerical [9], [10], [11] activity. Inspired by mean-field solutions of the spin-glass problem [12], [13], a simple picture for the dynamics was proposed in [14], based on the idea that metastable states act as traps in the phase-space with broadly distributed trapping times. This picture naturally leads to aging (i.e. non stationary dynamics) and suggests phenomenological laws for the decay of the thermoremanent magnetisation (and of the a.c. susceptibility) which are in quite good agreement with experimental data.

However, there are experimental features which are not compatible with the trap model proposed originally, which in fact corresponds to the phase space of the simplest kind – that of the ‘Random Energy Model’ (REM), for which Parisi’s $q(x)$ order parameter is simply $q(x < \bar{x}) = 0$ and $q(x > \bar{x}) = 1$ [15]. In that model, a metastable state is a single configuration; hence all dynamics is frozen if the system cannot “hop out”. Stated differently, there is no ‘bottom of the traps’ dynamics (see Fig 1-a).

This has various unsatisfactory consequences. For example, the equilibrium a.c. susceptibility is zero for all frequencies in this model, in plain contradiction with experiments. This can easily be cured by allowing fast, small scale fluctuations to reduce the ‘self-overlap’ $q(x > \bar{x})$ from 1 to a smaller value q_{EA} (the Edwards-Anderson order parameter). This ‘dressed’ REM behaviour was found recently in simple models for glasses [16], where the ‘traps’ can be very clearly identified, in particular in numerical simulations [17].

More importantly, a REM landscape is insufficient to account for the subtle effects induced by small temperature cycling [18]. Let us mention in particular the striking memory effect observed on the imaginary part of the a.c. susceptibility when the temperature is cycled as $T \longrightarrow T - \Delta T \longrightarrow T$. The signal rises strongly when the temperature is first decreased, showing that new dynamical processes are restarted. However, when the tem-

perature is raised again, the signal recovers exactly the value it had before the period at $T - \Delta T$, as if this period had not existed. A possible interpretation [18] is that, in our language, there are ‘traps within traps’ (see Fig. 1-b): the intermediate period corresponds to non-equilibrium dynamics within a trap – with hops between ‘supertraps’ frozen out.

The aim of this paper is thus to analyse in detail and generalize the REM-like trap model of ref. [14] (corresponding to a ‘one-step’ replica symmetry breaking (RSB) scheme [15]) to a fully foliated tree structure (‘full replica symmetry breaking’ [12]). A large amount of papers already studied various types of dynamics on trees [19], directly inspired from Parisi’s ultrametric construction. We however believe that our model is closer both to Parisi’s construction and to reality (in particular because most of these studies analyse deterministic trees). As in its REM version, aging appears most naturally within this framework and provides an excellent fit of experimental data, which in turn allow to determine the structure of the $q(x)$ characterizing the tree. We show that the data is already very well accounted for within a ‘two-step’ RSB approximation scheme. We discuss qualitatively T –jump experiments and ‘noise second spectrum’ [20] within our model. We speculate on how this ‘tree of states’ could be interpreted in finite dimensional space, and rephrase the droplet model of Fisher and Huse [21] in that context. We suggest that there is a spin-glass transition temperature associated with each length scale. We believe that our model, although still phenomenological, is of help to grasp the subtleties involved in the non-equilibrium dynamics of random or glassy systems. It also sheds light on the analytical results recently obtained for ‘toy’ microscopic spin-glass models exhibiting replica symmetry breaking [7], [6].

II. THE MODEL AND SOME ANALYTIC RESULTS

A. The Single Layer Tree (REM)

The simplest version of the model, already considered in [14], is defined as follows: the N possible ‘states’ of the system live on the leaves of the one layer tree drawn in Fig. 1-a.

To each branch α is associated an energy barrier ϵ_α distributed as (minus) the energy in the REM, i.e.:

$$\rho(\epsilon) = \frac{1}{T_g} \exp\left[-\frac{\epsilon}{T_g}\right] \quad (1)$$

where T_g is the glass transition temperature. To each α thus corresponds an Arrhenius hopping time $\tau_\alpha \equiv \tau_0 \exp \frac{\epsilon_\alpha}{T}$, where τ_0 is a microscopic time scale. From Eq. (1), one finds that the τ_α are (quenched) random variables distributed according to a power-law of index $x = \frac{T}{T_g}$:

$$p(\tau) = \frac{x\tau_0^x}{\tau^{1+x}} \quad (\tau \geq \tau_0) \quad (2)$$

We shall call τ an $[x]$ -variable. For $0 < x \leq 1$ (i.e. for $T < T_g$), the mean value of an $[x]$ -variable is infinite: this will lead to the aging phenomenon [14].

The model is further defined by saying that hopping rate from state α to state β is $\frac{1}{N\tau_\alpha}$. This simple dynamics has the virtue of giving the correct Boltzmann equilibrium distribution : for finite N , the equilibrium weight is: $P_{\text{eq}}^\alpha = Z^{-1} \exp \frac{\epsilon_\alpha}{T}$. The overlap between two states α, β is $q_0 \equiv 0$, while the self-overlap is taken to be $q_{EA} \leq 1$, independently of α . The ‘spin-spin’ correlation function $C(t_w + t, t_w) \equiv \langle S(t + t_w) S(t_w) \rangle$ within this model is thus naturally defined as:

$$C(t_w + t, t_w) \equiv q_{EA} \Pi_1(t, t_w) \quad (3)$$

where $\Pi_1(t, t_w)$ is the probability that the process has not jumped between t_w and $t_w + t$: this is the basic object we shall proceed to calculate.

Note that from Eq. (3), $C(t_w, t_w)$ can be smaller than 1. Strictly speaking, this is true in the limit $\tau_0 = 0$: it means that there is some dynamics taking place at short times, and reflecting the fact that a ‘state’ α is a ensemble of configurations mutually accessible within microscopic time scales. One could thus add branches connecting all these configurations to the ‘node’ α (see Fig 1-a); the ‘trapping’ times associated with this level of the tree are however such that their average is finite. A natural way is to continue Parisi’s $q(x)$ function for $x > 1$ and $q_{EA} < q < 1$ [22], such that the corresponding $\langle \tau \rangle = \frac{x}{1-x} \tau_0$ are microscopic. This corresponds to the *equilibrium* part of dynamics (see [23]), while $C(t, t_w)$ corresponds to

the non-stationary (aging) part. This important idea will be generalized below: for a finite number N of branches, or an upper cut-off in the distribution (2), the distinction between equilibrium and non-equilibrium becomes itself time-dependent.

The system is assumed to start in a randomly chosen initial metastable state. Rather than work at fixed t_w , it is convenient to work in Laplace transform, and define $\hat{\Pi}_1(t, E) = \int_0^\infty E dt_w \exp(-Et_w) \Pi_1(t, t_w)$. This is because, conditional on the value τ_α , the amount of time spent in a trap is exponentially distributed – using some standard properties of exponential random variables then makes calculations relatively straightforward.

We define $p(E, \tau_\alpha)$ to be the probability that at a random exponential time $\hat{t}_w$ of rate E the system is found in trap α with hopping rate $\frac{1}{\tau_\alpha}$. To be in trap α at time $\hat{t}_w$, one possibility is that the system starts in trap α and time $\hat{t}_w$ occurs before the system escapes from that trap. Clearly one starts in trap α with probability $\frac{1}{N}$, the probability that $\hat{t}_w$ occurs before escaping trap α is then given by property (3) of exponential random times as $E/(E + \frac{1}{\tau_\alpha})$. However, to start with, the system may be in any of the traps β $1 \leq \beta \leq N$ but escape from it before time $\hat{t}_w$. Conditional on this having happened, the memoryless property of the exponential time (property (A-1)) means that the probability of being in trap α at time $\hat{t}_w$ is just $p(E, \tau_\alpha)$, that is the game starts anew. Putting all of this together allows us to write down the renewal equation

$$p(E, \tau_\alpha) = \frac{1}{N} \frac{E\tau_\alpha}{E\tau_\alpha + 1} + \frac{1}{N} \sum_{\beta}^N \frac{1}{E\tau_\beta + 1} p(E, \tau_\alpha) \quad (4)$$

hence

$$p(E, \tau_\alpha) = \frac{\frac{E\tau_\alpha}{E\tau_\alpha + 1}}{\sum_{\beta}^N \frac{E\tau_\beta}{E\tau_\beta + 1}} \quad (5)$$

In the case where N is large we may use the approximation

$$p(E, \tau_\alpha) = \frac{\frac{E\tau_\alpha}{E\tau_\alpha + 1}}{N \langle \frac{E\tau_\beta}{E\tau_\beta + 1} \rangle} \quad (6)$$

the angled brackets indicating averaging over the disorder. In the regime where $E\tau_0 \ll 1$ (equivalent to $t_w \gg \tau_0$) we obtain

$$\langle \frac{E\tau_\beta}{E\tau_\beta + 1} \rangle \sim x\tau_0^x E^x c(x) \quad (7)$$

where

$$c(x) = \Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}$$

We now wish to calculate the probability that after an initial waiting time t_w the system has not decayed from the state it was in at time t_w after a subsequent time t has elapsed after t_w . Keeping the waiting time exponential, we find

$$\hat{\Pi}_1(t, E) = \left\langle \sum_{\beta}^N p(E, \tau_{\beta}) e^{-\frac{t}{\tau_{\beta}}} \right\rangle \sim \frac{1}{E^x c(x)} \int_{\tau_0}^{\infty} d\tau \tau^{-x-1} \frac{E\tau}{E\tau + 1} e^{-\frac{t}{\tau}} \quad (8)$$

Now, making a judicious change of variables and noting that $\exp(-Ew) \equiv \int_0^{\infty} E dv \exp(-Ev) \theta(v - w)$, one finally finds:

$$\Pi_1(t, t_w) = \frac{\sin(\pi x)}{\pi} \int_{\frac{t}{t+t_w}}^1 du (1-u)^{x-1} u^{-x} \quad (9)$$

This last form shows very clearly that the correlation function only depends, within this model, on the rescaled time $\frac{t}{t+t_w}$, i.e t over the ‘age’ of the system $t+t_w$: as argued in [14], only traps such that $\tau \sim O(t_w)$ have an appreciable probability to be observed after time t_w . From the same argument, the average energy decreases, in this model, as $\mathcal{E}(t_w) - \mathcal{E}(t=0) \sim -T \log(\frac{t_w}{\tau_0})$.

Two asymptotic regimes are of interest. If $t \ll t_w$ one obtains

$$\Pi_1(t, t_w) \sim 1 - \frac{\sin(\pi x)}{\pi(1-x)} \left(\frac{t}{t+t_w} \right)^{1-x} \quad (10-a)$$

In the regime $t \gg t_w$, we obtain

$$\Pi_1(t, t_w) \sim \frac{\sin(\pi x)}{\pi x} \left(\frac{t_w}{t+t_w} \right)^x \quad (10-b)$$

$C(t_w + t, t_w)$ has thus the ‘weak-ergodicity breaking’ property [14], [5]:

$$\lim_{t \rightarrow \infty} C(t_w + t, t_w) = 0 \quad \text{for all finite } t_w \quad (11-a)$$

but

$$\lim_{t_w \rightarrow \infty} C(t_w + t, t_w) = q_{EA} \quad \text{for all finite } t \quad (11-b)$$

meaning that ‘true’ ergodicity breaking only sets in after infinite waiting times [24].

The critical case $x = 1$ is of special interest, since it leads to aging but with an ‘anomalous’ scaling (i.e. not as $\frac{t}{t_w}$), we find:

$$\Pi_1(t, t_w) = \tau_0 \int_0^{\frac{t_w}{\tau_0}} du \frac{1 - \exp\left(-\frac{t_w + t - u\tau_0}{\tau_0}\right)}{(\log u)(t_w + t - u\tau_0)} \quad (12 - a)$$

which leads to

$$\Pi_1(t, t_w) = 1 - \frac{\log(\frac{t}{\tau_0})}{\log(\frac{t_w}{\tau_0})} \quad \text{for } \log(\frac{t}{\tau_0}) \ll \log(\frac{t_w}{\tau_0}) \quad (12 - b)$$

and

$$\Pi_1(t, t_w) = \frac{t_w}{t \log(\frac{t_w}{\tau_0})} \quad \text{for } t \gg t_w \quad (12 - c)$$

The above results pertain to the ‘aging’ case $x \leq 1$. For completeness, and also because they may be relevant to describe the equilibrium dynamics at short times (or for $T > T_g$, see [25], [22]), let us give the corresponding results for $x > 1$ in the limit $t_w \gg \tau_0$:

$$\Pi_1(t, t_w) = 1 - \frac{x}{x-1} \left(\frac{t}{\tau_0}\right) \quad \text{for } t \ll \tau_0 \quad (13 - a)$$

$$\Pi_1(t, t_w) = (x-1)\Gamma(x)\left(\frac{\tau_0}{t}\right)^{x-1} \quad \text{for } \tau_0 \ll t \ll t_w \quad (13 - b)$$

and

$$\Pi_1(t, t_w) = (x-1)^2\Gamma(x)\left(\frac{\tau_0^{x-1}t_w}{t^x}\right) \quad \text{for } t \gg t_w \quad (13 - c)$$

Note the major difference between these last results and those given in Eq. (10): for $\tau_0 \rightarrow 0$, $C(t_w + t, t_w)$ is identically zero for $x > 1$, but *remains non zero* for $x < 1$. In this sense, the crossing of $x = 1$ corresponds to a true dynamical freezing transition.

B. The Multi-Layer Tree (RSB)

The above single layer tree model can be generalized, following the construction of Mézard, Parisi and Virasoro for the tree of states (see [12], and also [26]). The hopping rate from state α to state β is set by a random escape time $\tau_{\alpha,\beta}$ the statistics of which depends on the overlap $q_{\alpha,\beta}$ between states. In Parisi’s tree, the free-energies associated to a certain level are distributed according to Eq. (1), but with a level-dependent T_g , and hence

with the overlap associated with that level. $\tau_{\alpha,\beta}$ is thus an $[x]$ -variable, with a q -dependent x which is precisely the inverse of Parisi's order parameter function $q(x)$. In the case of the REM ($q(x < \bar{x}) = 0$ and $q(x > \bar{x}) = q_{EA}$) we recover the single-layer tree since all the τ 's have the same statistics.

The procedure we follow to solve this model for a finite number M of layers (finite RSB) is a generalisation of the one presented above. We introduce the probabilities $\Pi_j(t, t_w)$, $j = 1, \dots, M$ that the process has never jumped beyond the j^{th} layer of the tree between t_w and $t_w + t$. $j = M$ corresponds to the deepest level of the tree and $j = 1$ the upper level of the tree – see Fig 1-b – and hence Π_1 is the same as the one introduced for the single level tree. Introducing the j^{th} level overlap q_j (with $q_0 = 0$ and $q_M = q_{EA}$), the spin-spin correlation function is clearly:

$$C(t_w + t, t_w) = \sum_{j=0}^M q_j [\Pi_j(t, t_w) - \Pi_{j+1}(t, t_w)] \quad (14)$$

with the convention that $\Pi_0(t, t_w) = 1$ and $\Pi_{M+1}(t, t_w) = 0$ (all processes inside a state have indeed taken place on the time scales $\gg \tau_0$ - see the discussion after Eq. (3).) The precise construction of the multi-layer tree is as follows, the $(j+1)$ th layer tree is constructed by adding N_{j+1} branches at the end of each branch of the j layer tree – the new states are now at the ends of these new branches. Associated with each new branch α is a new hopping rate $\frac{1}{\tau_{\alpha(j+1)}}$ with which the process hops back to level j and then falls back into the $(j+1)$ th layer. For simplicity, we define this hopping to be independent of all other hopping on the tree, i.e. the hopping dynamics is in parallel. We therefore have transitions from states characterised by inverse hopping rates $\{\tau_1, \tau_2, \dots, \tau_{k-1}, \tau_k, \dots, \tau_{j-1}, \tau_j\}$ to states $\{\tau_1, \tau_2, \dots, \tau_{k-1}, \tau'_k, \dots, \tau'_{j-1}, \tau'_j\}$ at rate $\frac{1}{\tau_k}$ (the primes denoting new rates chosen within the tree structure) and all these transitions are in parallel. The τ_j are quenched random variables chosen according to the law

$$p(\tau_j) = \frac{x_j \tau_{0j}^{x_j}}{\tau_j^{1+x_j}} \quad (\tau_j \geq \tau_{0j}) \quad (2')$$

where $0 < x_1 < x_2 \dots x_{j-1} < x_j < 1$. These times τ_j corresponds to (free) energy barriers ϵ_j distributed as in Eq. (1), but with a level dependent freezing temperature, T_{gj} . Hence,

again, the dynamics of this model reproduces, at long times and for a finite number of branches, the Boltzmann equilibrium: the total weight of a branch at level j is given by $Z_j^{-1} \exp \frac{\epsilon_j}{T}$.

For technical reasons we shall also assume that microscopic time scales associated with levels lower down the tree are much smaller than those associated with microscopic time scales further up the tree – this is also a reasonable hypothesis from the physical point of view. Therefore we shall impose $\tau_{0j+1} \ll \tau_{0j}$.

Once again we work with an exponential random time $\hat{t}_w$ of rate E . We define $p_j(E; \tau_1, \dots, \tau_j)$ to be the probability of finding the system in a set of traps characterised by $\{\tau_1, \dots, \tau_j\}$ at time $\hat{t}_w$. We calculate the p_j inductively as follows

$$p_{j+1}(E; \tau_1, \dots, \tau_{j+1}) = p_j(E; \tau_1, \dots, \tau_j) \cdot p(E_{j+1}, \tau_{j+1}) \quad (15)$$

where $E_j = E + \sum_{i=1}^{j-1} 1/\tau_i$ for $j \geq 2$, $E_1 = E$ and p is the result for the single layer tree. This comes from an application of Bayes' theorem and using properties (A-2) and (A-3) of exponential times. Consequently, $p_j(E; \tau_1, \dots, \tau_j) = \prod_{i=1}^j p(E_i, \tau_i)$. The fact that we have chosen $\tau_{0j} \ll \tau_{0(j-1)}$, and in addition chosen M to be reasonably small, means that we are assured that $E_j \tau_{0j} \ll 1$ and hence we may use the small E asymptotic result for the $p(E_j, \tau_j)$ in the expression above, yielding

$$p_j(E; \tau_1, \dots, \tau_j) = \prod_{i=1}^j \frac{\frac{E_i \tau_i}{E_i \tau_i + 1}}{N_i x_i c(x_i) E_i^{x_i} \tau_{0i}^{x_i}} \quad (16)$$

We now use

$$\hat{\Pi}_j(t, E) = \sum_{\{\tau_1, \dots, \tau_j\}} p_j(E; \tau_1, \dots, \tau_j) \exp(-t \sum_{i=1}^j \frac{1}{\tau_i}) \quad (17)$$

and the fact that the $N_i \rightarrow \infty$ and hence all combinations of $\{\tau_1, \dots, \tau_j\}$ exist, allows us to write

$$\hat{\Pi}_j(t, E) = \int_{\tau_{01}}^{\infty} d\tau_1 \cdots \int_{\tau_{0j}}^{\infty} d\tau_j \exp\left(-t \sum_{i=1}^j \frac{1}{\tau_i}\right) \prod_{i=1}^j \frac{\frac{E_i \tau_i^{-x_i}}{E_i \tau_i + 1}}{c(x_i) E_i^{x_i}} \quad (18)$$

One again attempts an integration by substitution, starting with τ_j and proceeding down to τ_1 . On inverting the Laplace transform one obtains

$$\Pi_j(t, t_w) = \frac{\int_0^1 \prod_{i=1}^j du_i u_i^{-x_i} (1 - u_i)^{x_i - 1} \theta(\prod_{i=1}^j u_i - \frac{t}{t+t_w})}{\prod_{i=1}^j B(x_i, 1 - x_i)} \quad (19)$$

where θ is the Heaviside step function. The above formula has an immediate probabilistic interpretation. If U_i are independent random variables from Beta distributions of indices $1 - x_i$ and x_i respectively, then

$$\Pi_j(t, t_w) = P\left(\prod_{i=1}^j U_i > \frac{t}{t + t_w}\right) \quad (20)$$

A useful rearrangement of equation (19) for asymptotic analysis is

$$\Pi_{j+1}(t, t_w) = \Pi_j(t, t_w) - \frac{\int_0^1 \prod_{i=1}^j du_i u_i^{-x_i} (1 - u_i)^{x_i-1} \theta(\prod_{i=1}^j u_i - s) B_{\frac{s}{\prod_{i=1}^j u_i}}(1 - x_{j+1}, x_{j+1})}{\prod_{i=1}^{j+1} B(x_i, 1 - x_i)} \quad (21)$$

where $s = \frac{t}{t + t_w}$ and

$$B_\gamma(x, y) \equiv \int_0^\gamma du u^{x-1} (1 - u)^{y-1} \quad (22)$$

is the incomplete Beta function (clearly $B_1(x, y) \equiv B(x, y)$). The above rearrangement also makes clear that the obvious inequality $\Pi_{j+1}(t, t_w) < \Pi_j(t, t_w)$ is satisfied.

Let us focus on the short time and long time asymptotics implied by the above analytical forms. For $t \ll t_w$, we find that

$$\begin{aligned} C(t_w + t, t_w) &= q_{EA} - (q_{EA} - q_{M-1}) \frac{\sin(\pi x_M)}{\pi(1 - x_M)} \prod_{j=1}^{M-1} \frac{B(x_j, x_M - x_j)}{B(x_j, 1 - x_j)} \left(\frac{t}{t + t_w}\right)^{1-x_M} + \\ &+ O\left(\left(\frac{t}{t_w}\right)^{1-x_{M-1}}\right) \end{aligned} \quad (22)$$

This form shows that the initial decay is, as expected, dominated by the deepest level of the tree. The possibility of ‘jumping’ to higher levels during the waiting period however acts to reduce the effective waiting time by a factor $\left[\prod_{j=1}^{M-1} \frac{B(x_j, x_M - x_j)}{B(x_j, 1 - x_j)}\right]^{-\frac{1}{1-x_M}} < 1$. Each of these large scale jumps indeed completely restarts the small scale dynamics.

In the limit $t \gg t_w$, on the other hand, the asymptotic decay is given by:

$$C(t_w + t, t_w) = q_1 \frac{\sin(\pi x_1)}{\pi x_1} \left(\frac{t_w}{t + t_w}\right)^{x_1} + O\left(\left(\frac{t_w}{t}\right)^{x_1+x_2}\right) \quad (23)$$

Note that, again, $C(t_w + t, t_w)$ is only a function of the ratio $\frac{t}{t_w}$. This would not be true if the number of branches N_j is finite, as we shall discuss now.

C. Finite number of branches and interrupted aging.

Let us now discuss the case where the number of ‘states’ N is finite, starting with the single layer tree model. As can be seen from Eq. (5), or directly from the argument giving the order of magnitude of the largest τ drawn from distribution (1), the results established in section II.1 rely on the inequality:

$$t + t_w \ll \tau_0 N^{\frac{1}{x}} \equiv t_{\text{erg}} \quad (24)$$

In the opposite case, the dynamics ceases to depend on t, t_w , and stationary dynamics resume and aging is ‘interrupted’: t_{erg} is the ergodic time of the problem. Π_1 asymptotically behaves as in Eqs. (10), but with the variable $\frac{t}{t_w}$ replaced by $\frac{t}{t_{\text{erg}}}$. The scaling of $C(t_w + t, t_w)$ as a function of $\frac{t}{t_w}$ is thus expected to degrade progressively as t, t_w approach $t_{\text{erg}}(N)$. This was proposed as an explanation for the small (but systematic) deviation from a perfect $\frac{t}{t_w}$ scaling observed experimentally [14]. Let us note that very similar results would of course hold if N is infinite but the power-law distribution (2) happens to be truncated beyond a certain t_{erg} .

The case of a multilayer tree is interesting, but rather complex – we shall thus only give qualitative arguments. Let us assume that the ergodic time at level j is now t_{erg}^j . Is it reasonable to assume that the hierarchy of the x_j (slower dynamics as one gets higher up the tree) is maintained for the ergodic times, i.e.: $t_{\text{erg}}^j \gg t_{\text{erg}}^{j+1}$. For a given waiting time t_w , a specific level of the tree j_w is naturally selected through the condition

$$t_{\text{erg}}^{j(t_w)} \gg t_w \gg t_{\text{erg}}^{j(t_w)+1} \quad (25)$$

meaning that all the levels ‘below’ $j(t_w)$ are equilibrated, while those above are still aging. In this case, even if $x_{j(t_w)+1} < 1$, the corresponding transitions are equilibrated – reducing further the effective value of q_{EA} , renormalized by small scales dynamics. The interesting point, however, is that the effective ‘ergodic’ time for which deviations from a $\frac{t}{t_w}$ scaling first become visible is now $t_{\text{erg}}^{j(t_w)}$ which becomes, through Eq. (24), *dependent on t_w itself*.

[It might well be that the ergodic time determined experimentally in [14] is only the one corresponding to the particular level of the tree which happens to be probed on the $1 - 10000$ seconds time scales].

Another consequence of these nested equilibrium time scales is the following: if a perturbation is applied to the system at time t_w , the response at time $t_w + t$ will depend on how much the system has been able to reequilibrate during time t . The response is thus expected to depend on the particular level j defined as in Eq. (25), but with t replacing t_w . Hence, for $t \ll t_w$, one expects that $C(t_w + t, t_w) = q_{j(t)}$, leading to violations of the $\frac{t}{t_w}$ scaling at short times too. Said differently, the effective ‘microscopic’ time scale τ_0 is replaced by $t_{\text{erg}}^{j(t_w)+1}$ [27]. The existence of different time domains in the plane t_w, t was in fact suggested in refs. [6], [7].

III. RELATION WITH MEASURABLE QUANTITIES.

A. Constant temperature data

Although noise measurements can be and have been performed on spin-glasses to access directly the correlation function computed above, the response function – measured through the thermoremanent magnetisation (TRM) or the a.c. susceptibility – is of much easier access. Most of the available data thus gives information on the response function $R(t, t') \equiv \frac{\partial \langle S(t) \rangle}{\partial H(t')}$ (but see below). For equilibrium dynamics, the response and correlation functions are related through the Fluctuation-Dissipation theorem (FDT). For non-stationary dynamics, an extended form of this theorem has recently been proposed [5], [7], [6], [8]. This relation can be written as:

$$T R(t, t') = \theta(t - t') X[C(t, t')] \frac{\partial C(t, t')}{\partial t'} \quad (26 - a)$$

where $X(\cdot)$ is a certain function which can be computed within simple models [5], [7], [6]. For the SK model near T_g and the toy-model of [6], X is the inverse of the function $q(x)$, whereas for the spherical p -spin model, X is a constant ≤ 1 (depending on temperature).

(Equilibrium dynamics corresponds to $X \equiv 1$ and one recovers the usual FDT.)

Within the present context, the response function can be computed by assuming that to each trap i is associated a certain magnetisation m_i with probability $\rho(m)$, which we shall take to be even $\rho(-m) = \rho(m)$. The hopping rate from state i to state j is modified in the presence of a field H by a factor $\exp - \left[\frac{H\{\zeta m_i - (1-\zeta)m_j\}}{T} \right]$ which ensures the correct equilibrium weighting for all values of ζ . It is then easy to establish that [28]:

$$T R(t, t') = \theta(t - t') \left[(1 - \zeta) \frac{\partial C(t, t')}{\partial t'} - \zeta \frac{\partial C(t, t')}{\partial t} \right] \quad (26 - b)$$

If $C(t, t') = C(t - t')$, the usual FDT is recovered. If – as is the case here – $C(t, t') = C(\frac{t}{t'})$, then one finds that Eq. (26-a) holds with $X = (1 - \zeta) + \zeta \frac{t'}{t} \equiv (1 - \zeta) + \frac{\zeta}{C^{-1}(\frac{t}{t'})}$. Note that in the early regime corresponding to equilibrium dynamics ($t = t' + O(\tau_0)$), $X \equiv 1$ independently of ζ .

Now, the TRM $M(t_w + t, t_w)$ is defined as:

$$M(t_w + t, t_w) = \int_0^{t_w} dt' R(t_w + t, t') H \quad (27)$$

where H is the applied field between 0 and t_w . Defining the function $Y(q)$ through $X(q) = \frac{dY(q)}{dq}$ and using Eq. (26), we obtain that the experimentally measured TRM and $C(t_w + t, t_w)$ are related through [5]:

$$M(t_w + t, t_w) = Y[C(t_w + t, t_w)] h \quad (28)$$

where $h \equiv \frac{H}{T}$. The simplest assumption of a REM with $q(x < \bar{x}) = 0$, $q(x > \bar{x}) = q_{EA}$ and $X = \text{const.}$ leads to:

$$M(t_w + t, t_w) = q_{EA} X h \Pi_1(t, t_w) \quad (29)$$

Experimentally, the decay of $M(t_w + t, t_w)$ at short and large times is indeed very well described by the forms given in Eqs. (10-a, 10-b). In particular, the scaling as a function of the reduced time $\frac{t}{t_w}$ is reasonably well obeyed (but see [14]-b) and section II.3 above). In fact, the experimental value of $\bar{x}$ is in the range 0.6 – 0.9 if extracted using the short-time expansion (10-a), and in the range 0.05 – 0.2 if extracted using the long-time expansion (10-b) [14], where it was called γ . This suggests that a more complicated RSB scheme is needed,

generating a tree with at least two levels – allowing for the existence of two independent exponents x_1 and x_2 . In the spirit of Parisi’s step by step construction of the SK solution, we shall thus assume that $q(x)$ is given by:

$$q(x < x_1) = 0 \quad q(x_1 < x < x_2) = q \quad q(x > x_2) = q_{EA} \quad (30)$$

In order to obtain $M(t_w + t, t_w)$ we first assume that $X = \text{const.}$, or $\zeta = 0$. This assumption is natural within our picture since q_j corresponds to the fraction of ‘frozen’ spins at level $j - 1$: these spins thus retain their magnetisation when the process ‘jumps’ at level $j - 1$ leading to:

$$M(t_w + t, t_w) \propto [(q_{EA} - q)\Pi_2(t, t_w) + q\Pi_1(t, t_w)] \quad (31)$$

This form is only valid after the short time ($x > 1$) relaxation has taken place; just before the field is cut, the field-cooled magnetisation is by definition $M(t_w^-, t_w) := \chi_{FC}H$, where χ_{FC} is roughly temperature independent. We choose units such that $M(t_w^-, t_w) \equiv 1$. The four fitting parameters are thus $x_1, x_2, \frac{q}{q_{EA}}$ for the *shape* of the TRM decay and $\frac{q_{EA}X}{T}$ for the *y-axis* scale. We show in Fig 2-a and 2-b the best fits obtained for both studied samples, Ag:Mn_{2.6%} (AgMn) at $T = 9K$ ($0.87 T_g$) and CdCr_{1.7}In_{0.3}S₄ (CrIn) at $T = 10K$ ($0.6T_g$), plotted as a function of the rescaled variable $s = \frac{t}{t+t_w}$. As can be seen, these fits are excellent on the whole time regime, which extends, for CrIn, from 10 sec. to 2.3×10^5 sec [29]. The value of the fitting parameters for different temperatures and samples are given in Table 1 and 2. Assuming that X is not critical at T_g , we can extract values for the exponents β, β' defined as $q_{EA} \simeq (\frac{T-T_g}{T_g})^\beta$ and $q \simeq (\frac{T-T_g}{T_g})^{\beta'}$: $\beta \simeq 0.65$ and $\beta' \simeq 1.0$ (both samples give the same values for these exponents, within 10%.) Our value of β is compatible with other determinations ($\beta \simeq 0.5 - 1$) [30].

The fits shown in Fig. 2-a, 2-b implicitly assume that $x_2 < 1$. We have also tried the case $x_2 = 1$ – the free parameter now being τ_0 . The resulting fit is of much poorer quality; however, as will be discussed in section IV (Fig 6 below), the initial part of the TRM decay for different t_w is rather well rescaled when plotted as a function of $\frac{\log(\frac{t}{\tau_0})}{\log(\frac{t_w}{\tau_0})}$, as suggested by Eq. (12-b). We interpret this as a sign that $q(x)$ is in fact continuous, at least near $x = 1$.

We have also tried to fit the data using different values of ζ . The quality of the fit remains excellent if $\zeta \leq \frac{1}{2}$ (although the values of q_{EA}, q are somewhat changed), and deteriorates for larger values of ζ . One can also use the form of $X(C)$ suggested by the models studied in [5], [7], [6], i.e., the inverse of the function $q(x)$ defined by Eq. (30). Defining a characteristic time t^* such that $C(t_w + t^*, t_w) = q$, the TRM takes the following form (for $t = 0^+$):

$$M(t_w + t, t_w) = [x_1(C(t_w + t, t_w) - q) + x_2 q]h \quad (t < t^*) \quad (32 - a)$$

$$M(t_w + t, t_w) = x_2 C(t_w + t, t_w)h \quad (t > t^*) \quad (32 - b)$$

i.e., the decay rate is discontinuous at $t = t^*$. We have plotted in Fig 3 the best fit obtained using Eqs (32): the initial and final part of the curve can be accounted for, but not the intermediate region around $t = t^*$. This suggests that either the function X is (in our case) not connected simply to $q(x)$ and is a constant for the range of correlations probed, or else that a continuous RSB scheme is needed (to remove any discontinuity in decay rate). More work on this point is certainly needed, in particular to extend our formulae to a continuous tree.

Let us now turn to the a.c. susceptibility measurements. The object which is measured is now $\chi(\omega, t) = \int_{-\infty}^t R(t, t') \exp(i\omega(t' - t)) dt'$. Let us first assume a REM-like dynamics. Then using Eqs. (26), and in the limit $t \gg \omega^{-1}$ (actually needed for an accurate measurement of $\chi(\omega, t)$), we find that:

$$\chi(\omega, t) = \frac{X q_{EA}}{T} \int_0^\infty du (\exp[-i(\omega t)u] - 1) \frac{\sin(\pi \bar{x})}{\pi} u^{-\bar{x}} \quad (33 - a)$$

(the $\omega = 0$ value of χ' has been removed). Defining $\bar{\omega} \equiv \omega t$ and changing variables,

$$\chi(\omega, t) \equiv \chi(\bar{\omega}) = X \frac{q_{EA}}{T} \frac{\sin(\pi \bar{x})}{\pi} \Gamma(1 - \bar{x}) e^{-i\delta} \bar{\omega}^{\bar{x}-1} \quad (33 - b)$$

in agreement with the arguments given in [14]. Eq. (33-b) shows that:

(i) the relevant variable is again the rescaled time ωt : this is very well obeyed by experimental data.

(ii) There is a ‘constant loss angle’ $\delta = \frac{\pi}{2}(1 - \bar{x})$ between the aging part of χ' and χ'' , as expected from Kramers-Kronig relations.

(iii) For a multilayer tree, the same formula (33-b) will hold in the limit $\bar{\omega} \gg 1$, with $\bar{x}$ replaced by x_M . In other words, the a.c. susceptibility measurements are only sensitive to the deepest level of the tree. Actually, Eq. (33-b) for χ'' is very well obeyed experimentally – although a ‘non aging’ contribution χ''_{eq} must be added: see Fig. 4. Indeed, in the limit $\bar{\omega} \rightarrow \infty$, the short time dynamics corresponding to intra-state fluctuations ($x > 1$) cannot be neglected in Eq. (33).

As mentioned above, magnetic noise measurements have also been performed in order to test FDT in spin-glasses [31]. The ‘usual’ FDT would predict that the noise spectrum $S(\omega, t)$ is related to $\chi''(\omega, t)$ through:

$$\chi''(\omega, t) = \frac{\pi X}{2} \frac{\omega S(\omega, t)}{T} \quad (34)$$

with $X \equiv 1$. The non-equilibrium theory proposed in [5], [7], [6] claims that $X = x_M < 1$ in the limit $\omega t \gg 1$, while our model suggests (in the same limit) $X = 1 - \frac{\zeta}{\omega t}$ – see also [32]. Experimentally, the noise measurements were not calibrated in an absolute way. The proportionality relation (34) was indeed confirmed but no value of X could be determined. In view of the present discussion, it seems to us that this would be a very important point to check: the value of x_M extracted from the decay of χ'' or the TRM should be the same as the proportionality constant extracted from Eq. (34). Since x_M is in the range $0.6 - 0.9$, a significant effect could show up by comparing carefully equilibrium and non-equilibrium regimes.

Let us finally note that for $x > 1$, $\chi''(\omega, t)$ is given by:

$$\chi''(\omega, t) \propto \omega \tau_0^{x-1} t^{2-x} \quad \text{for } \omega \ll t^{-1} \quad (35 - a)$$

$$\chi''(\omega, t) \propto (\omega \tau_0)^{x-1} \quad \text{for } t^{-1} \ll \omega \ll \tau_0^{-1} \quad (35 - b)$$

(the large frequency region $\omega \gg \tau_0^{-1}$ requires a detailed description of $p(\tau)$ for $\tau \ll \tau_0$). Hence for $x \simeq 1$ (i.e slightly above the spin glass transition, or equilibrium dynamics in the spin-glass phase), Eqs (33-35) gives $S(\omega) \propto \frac{T}{\omega}$ i.e. the model naturally leads to $1/f$ noise.

B. Temperature cycling

As mentioned in the introduction, more sophisticated experimental protocols have been investigated [18], [33], where the temperature or magnetic field is changed *during* TRM or χ'' measurements. A complete analysis of these experiments is beyond the aim of the present paper and will be given in separate publications [34]. We shall restrict here to a qualitative discussion of the results, focusing on the remarkable memory effect mentioned in the introduction, and illustrated in Fig. 5 (from [35]).

Let us thus discuss the effect of a small temperature change on χ'' within a REM landscape. As discussed in the previous section, this REM description is sufficient to account for the behaviour of $\chi''(\omega, t)$ in the limit $\omega t \gg 1$ (provided an equilibrium contribution is added). In the REM, the (free-) energy landscape does not evolve with temperature and $\bar{x}$ is simply given by $\frac{T}{T_g}$. A small change of temperature $T \longrightarrow T' = T - \Delta T$ simply changes the trapping times as: $\tau \longrightarrow \tau' = \tau_0(\frac{\tau}{\tau_0})^p$ with $p = \frac{T}{T'} > 1$. Suppose that the waiting time before the small temperature jump is t_{w1} – see Fig 5. As emphasized in [14] and above, only traps with $\tau \simeq t_{w1}$ have a significant probability to be observed. Hence, if the time spent at temperature T' is such that $t_{w2} \ll \tau_0(\frac{t_{w1}}{\tau_0})^p$, it is quite obvious that no jump will take place during this intermediate period. The ‘memory effect’ $\chi''(\omega, t_{w1} + t_{w2} + 0) = \chi''(\omega, t_{w1} - 0)$ would then be trivial. On the other hand, one can show that the change of χ'' at t_{w1} is given by:

$$\frac{\chi''(\omega, t_{w1} + 0)}{\chi''(\omega, t_{w1} - 0)} = p \frac{q_{EA}(T')X(T')}{q_{EA}(T)X(T)} (\omega\tau_0)^{(\bar{x}-1)(\frac{1}{p}-1)} \quad (36)$$

Typically (see Fig 5), $p = 1.2$, $\bar{x} = .75$, while the variation of q_{EA} is roughly a factor 1.2. Hence, provided $\tau_0 < 10^{-4}$ sec., χ'' should first decrease and then remain constant when the temperature jumps down, at variance with the experimental results. In physical terms, all dynamical processes should be slowed down by the temperature jump – hence the decrease of χ'' and the memory effect. The *simultaneous* observation of a strong increase of χ'' and $|\frac{\partial \chi''}{\partial t}|$ (‘rejuvenation’) and of the memory effect is non trivial [36]. In [18], it was suggested that the origin of this combined effect comes from the hierarchical nature of the phase-

space, with valleys bifurcating into subvalleys at all temperatures below T_g in a continuous sequence of ‘micro phase transitions’. Even a small temperature change can thus be thought as a ‘quench’ from ‘high’ temperature, setting back the age of the system to zero – hence the increase of χ'' .

We want to rephrase this interpretation within the framework developed here. As mentioned above, the construction of Parisi’s tree can be extended ‘within states’ to $x > 1$ and $q > q_{EA}$, simply reflecting the fact that even within each state there are long-lived configurations and non exponential equilibrium dynamics, even above T_g . However, as mentioned previously, only the part of the tree corresponding to $x < 1$ corresponds to non-stationary dynamics, while the part $x > 1$ is equilibrated within microscopic time scales. T_g is the temperature at which the smallest x first reaches the critical value 1, but one can expect a whole sequence of phase transitions as the temperature is decreased, corresponding to the crossing of $x = 1$ for the successive levels of the tree, much as in the ‘generalized’ Random Energy Model [37]. (As discussed below, this could correspond in real space to the progressive freezing of the dynamics over smaller and smaller length scales.) Hence if at a certain temperature $x_M < 1$ but $x_{M+1} > 1$, only the transitions at the M^{th} level of the tree will contribute to the relaxation of χ'' . But if upon lowering the temperature x_{M+1} becomes less than 1, then the processes corresponding to this level of the tree will start contributing, while those corresponding to x_M are to a certain extent frozen – allowing for the memory effect. The small temperature jump is furthermore tantamount to a *quench* for the $M + 1$ level since when $x_{M+1} > 1$, all states are more or less equivalent, corresponding to a random initial condition at $t = t_{w1} + 0$.

Another related scenario is also possible: as discussed in section II.3, the finiteness of the ergodic time t_{erg}^j at level j leads to ‘interrupted aging’ [14], after a time t_{erg}^j . The crucial levels of the tree are in this case not those such that $x_M < 1 < x_{M+1}$ but, as discussed in section II.3, those such that $t_{\text{erg}}^M(T) \gg t_w \gg t_{\text{erg}}^{M+1}(T)$, even if $x_{M+1} < 1$. Aging processes will be restarted upon cooling if $t_{\text{erg}}^{M+1}(T - \Delta T) \gg t_w$. A quantitative analysis of the experimental data along these lines is currently underway. Experiments investigating

the role of the magnetic field on aging and their interpretation are reported in [27].

C. Noise second spectrum

Another interesting set of experiments which can be discussed within our model is the noise ‘second spectrum’ measurements of Weissmann et al. [20]. Very briefly, the point is that on sufficiently small samples, one observes $\frac{1}{f}$ noise, but with an amplitude which is itself ‘noisy’, i.e. randomly changing with time (albeit on rather long time scales.) The natural interpretation – discussed in different terms in [20] – is that the noise primarily comes from the near equilibrium levels $x \simeq 1$, but that higher level jumps will slightly change the amplitude of this noise – since the systems are small enough, the averaging over different ‘branches’ is not performed. For a multilayer tree, the prediction is obviously that the noise spectrum behaves as $\omega^{x_M-2} \simeq \omega^{-1}$, but with a *noisy amplitude* with a correlation function decaying as t^{-x_1} , as observed experimentally (where this exponent is called β). A more quantitative analysis of the experimental results along these lines would however be desirable, but we note that the value of $\beta = x_1$ determined for CuMn in [20] is very close to the one quoted in Table 1,2 for x_1 , and evolves similarly with temperature.

IV. DISCUSSION. TREES AND DROPLETS

Fisher and Huse [21], see also [38] were the first to stress the importance of understanding the nature of the spin-glass excitations in finite dimensional spaces. They argued that these excitations should be of the form of *compact* ‘droplets’, with a surface much smaller than their volume. These droplets were defined with respect to the (unique up to a global spin-flip) ground state as the ‘spin-flip’ excitation of lowest possible energy within a region of size L ; the energy of such a droplet is found to be of order L^θ ($\theta \simeq 0.2$ in $d = 3$), and the energy barrier for such a droplet to be activated grows as L^ψ . Apart from the presence of these ever-growing energy barriers, the description of Fisher and Huse of a spin-glass is that of a ‘disguised ferromagnet’. In particular, exactly as in a ferromagnet quenched in zero field,

a spin-glass quenched below T_g would approach equilibrium by growing domains of the two ‘pure phases’ assumed to be present, thereby eliminating the extra energy (L^θ) associated with the domain walls. The system thus *coarsens* with time; after time t_w the typical size of the domains is $R(t_w) = \log^{\frac{1}{\psi}}(\frac{t_w}{\tau_0})$ – instead of $R(t_w) = t_w^{1/2}$ in a ferromagnet (with non conserved dynamics, see [39]). Before discussing the form of the relaxation proposed in [21] inspired from this picture, it is useful to recall how $C(t_w + t, t_w)$ would look like for a simple ferromagnet. An exactly soluble case is the ϕ_n^4 theory in the limit $n \rightarrow \infty$, for which one finds [39]:

$$C_{n=\infty}(t_w + t, t_w) = \left(\frac{4(t_w + t)t_w}{(2t_w + t)^2} \right)^{\frac{d}{4}} \quad (37)$$

(d is the dimension of space). Mazenko’s approximate theory for $n = 1$ [8], [39] leads to:

$$C_{n=1}(t_w + t, t_w) = \frac{4}{\pi} \tan^{-1} \sqrt{\frac{2 - C_{n=\infty}(t_w + t, t_w)}{C_{n=\infty}(t_w + t, t_w)}} - 1 \quad (38)$$

For $t \gg t_w$, expressions (37,38) simplify to $(\frac{t_w}{t})^{d/4} \equiv (\frac{R(t_w)}{R(t)})^\lambda$ with $\lambda = \frac{d}{2}$, whereas more elaborate theories show that λ is non trivial in general [39]. Fisher and Huse argued that the appearance of the ratio of two length scales, $R(t)$ and $R(t_w)$, should be general and proposed for the spin-glass dynamics to write:

$$C_{FH}(t_w + t, t_w) \propto R(t)^{-\theta} \Sigma\left(\frac{R(t)}{R(t_w)}\right) \quad (39)$$

with $\Sigma(x \rightarrow 0) = \text{const.}$ and $\Sigma(x \rightarrow \infty) = x^{\theta-\lambda}$. (Again, we identify $C(t_w + t, t_w)$ and $M(t_w + t, t_w)$ in the small H limit). Eq. (39) means that:

- a) The scaling variable for aging should be $\frac{\log(\frac{t}{\tau_0})}{\log(\frac{t_w}{\tau_0})}$ rather than $\frac{t}{t_w}$.
- b) Even in the limit where $t_w = \infty$, $C_{FH}(t_w + t, t_w)$ decays to zero at large times if $\theta > 0$, at variance with Eq. (11-b) and the results of [5], [7], [6].

Before comparing with the experimental data, let us remark that, from Eqs. (37-38), $C(t_w + t, t_w)$ is a function of $\frac{t}{t_w}$ *over the whole range* of time scales. This scaling variable is thus more general than the asymptotic one, i.e. $\frac{R(t)}{R(t_w)}$. This suggests that even in the case of logarithmic domain growth, a $\frac{t}{t_w}$ scaling should hold for coarsening. We have confirmed

this [40] for the 1d Random Field Ising Model where $R(t) \propto \log^4 t$ (see e.g. [41], [42]). As emphasized in [8], Eqs. (37-38) show that aging is not specific of spin-glasses, but occur as soon as the equilibration time of the system is infinite (or much larger than t, t_w). We have plotted in Fig. 6 $C_n(t_w + t, t_w)$ as given by Eqs. (37-38) as a function of $\frac{t}{t_w + t}$. The major difference with the spin-glass data is the singular behaviour of the latter for short times (see Figs. 2-a,b).

We have tried to test the predictions contained in Eq.(39). For example, we have looked for the best values of θ and τ_0 to rescale TRMs at a fixed temperature and different t_w , for both AgMn and CrIn. The data points unambiguously towards $\theta = 0$; data collapse is quite good at short times but deteriorates as soon as $t > t_w$: see Fig. 7. The other test is to compare different temperatures with the same t_w . We fixed $\theta = 0$, and chose $\tau_0(T)$ and the T-dependent proportionality constant in Eq. (39) (which is essentially $q_{EA}(T)$) to best rescale the *late* part of the curves. Although this late part can indeed be satisfactorily fitted by $\log^{-\frac{\lambda}{\psi}}(\frac{t}{\tau_0(T)})$ with $\frac{\lambda}{\psi} \simeq 1.2$, the early part of the data does not scale (Fig. 8). Hence we believe that Eq. (39), even with $\theta = 0$, is inadequate to describe the data consistently.

Koper and Hilhorst [38] proposed a similar picture, although they assumed that $R(t)$ grows as a power-law t^p rather than logarithmically. Also, the relaxation time associated to a domain of size R is taken to be $\propto R^z$. It would be too long to discuss in detail this theory here; let us simply mention that it does lead a relaxation of $\chi''(\omega, t)$ behaving as $\omega^{-1}t^{-pz}$. Consistency with the ωt (or $\frac{t}{t_w}$) scaling observed experimentally leads to $pz \simeq 1$ – which is in fact natural since it means that the growth time and relaxation time of droplets are comparable [43]. However, the resulting relaxation of χ'' as t^{-1} is much too fast (compare with Eq. (34), with $\bar{x} \simeq .75$).

The basic remark of Fisher, Huse, Koper and Hilhorst that the flipping spins are clustered somewhere in space and that the time scales should grow with the size of these clusters seems however unavoidable. How can this be reconciled with replica symmetry breaking ? Of course, a proper replica (or dynamical) theory in finite dimension is needed to answer

completely this question. Such a theory is not yet available for spin glasses, but has been worked out for the simpler problems of manifolds in random media [44], [45] (i.e. polymers, surfaces, vortex lattices. etc..). One basically finds that $q(x)$ becomes L (scale) dependent, with a characteristic value of $x(L)$ varying as $L^{-\theta}$: small scales correspond to large x .

From Eq. (1) and the interpretation of $x(L)$, one thus sees that the energy distribution for the excitations ('droplets') of scale L has the form:

$$\rho(f, L) = \frac{1}{f_0 L^\theta} \exp\left(-\frac{|f|}{f_0 L^\theta}\right) \quad (40)$$

showing, as postulated by Fisher and Huse, that the energy scale grows as L^θ . At this stage, an important difference with Fisher and Huse is the exponential form of the distribution. Interestingly, Eq. (40) means, within our interpretation, that there is a spin-glass transition temperature $T_g(L)$ associated with each scale, defined through $x(L, T_g(L)) = 1$. The infinite sequence of micro-phase transitions suggested in [3], [18] thus corresponds to a progressive 'weak ergodicity breaking' (in the sense that $x(L)$ crosses 1, see II.1) of smaller and smaller length scales (faster and faster degrees of freedom).

Quite naturally, one can associate an L -dependent overlap $q(L)$ between configurations obtained by flipping a droplet as $1 - q(L) \propto (\frac{L}{\xi})^{d_f}$, where d_f is the fractal dimension of the droplets (d_f is equal to d for Fisher and Huse) and ξ is a correlation length. Hence, the picture we propose is in fact very close the original droplet model, except that:

- a) The 'droplets' are only required to be metastable, and not 'lowest' excitations,
- b) These droplets may a priori be non compact ($d_f < d$), and
- c) The time scale associated with droplets of size L is *not peaked* around some $\tau(L)$, but rather *power-law distributed* with a parameter $x(L)$ which becomes smaller and smaller at large sizes.
- d) The Fisher-Huse time scale $\tau_0 \exp(L^\theta)$ could be interpreted as the ergodic time $t_{\text{erg}}(L)$ associated to scale L : one indeed should expect that Eq. (40) ceases to be valid for $|f| \gg f_0 L^\theta$.
- e) Aging will be totally interrupted when the 'terminal' ergodic time scale $t_{\text{erg}}(\xi)$ associ-

ated to ξ is reached.

This picture suggests that the final description of the experiments should involve a continuous tree, rather than the 2-level approximation that we have chosen. Further work on this aspect and on the related problem of the Fluctuation Dissipation theorem is certainly needed; we however hope that the scenario proposed here can be helpful to think about non ergodic dynamics in glassy systems, in particular those in which quenched disorder is a priori absent [16]. We note in this respect that the spontaneous appearance of power-law distributed ‘trapping’ times with x crossing 1 at the glass transition has been reported for hard-sphere systems [46].

Acknowledgments We wish to express warm gratitude towards A. Barrat, L. Cugliandolo, S. Franz, J. Hammann, M. Mézard, M. Ocio, E. Vincent, M. Weissmann for enlightening critical discussions. We also take this opportunity to acknowledge the inspiring work of L. Cugliandolo, S. Franz, J. Kurchan and M. Mézard on this problem.

Figure Captions

Figure 1 Schematic views of the free energy landscape with associated Parisi trees. 1-a: REM landscape, 1-b: Full RSB landscape.

Figure 2 2-a : Fit of TRM decay for CrIn at 10 K against two level tree theory with simple FDT. Note that the fit is perfect over the whole time domain. 2-b : Fit of TRM decay for AgMn at 9 K against two level tree theory with simple FDT.

Figure 3 Fit of TRM decay for AgMn at 9 K against two level tree theory with generalized FDT as given by equation (26).

Figure 4 Fit of the decay of the out of phase susceptibility of CrIn (at T=12 K, $\omega = 10^{-2}$ Hz.) as $\chi''(\omega, t) - \chi_{\text{eq.}}(\omega) \propto t^{x_2-1}$, and $x_2 = .79$ (see Table 2).

Figure 5 Sketch of the temperature cycling protocole and the evolution of χ'' during this cycling (from [35]). Note the *strong spike* just after the temperature decrease followed by a *perfect ‘memory’* in the third stage.

Figure 6 Aging in $n = 1$ (approximate result) and $n = \infty$ (exact result) ferromagnets.

Note that the small time behaviour is regular, at variance with the experimental data on spin glasses (Figs 2-a, 2-b).

Figure 7 $M(t_w + t, t_w) \log^{\theta/\psi}(\frac{t}{\tau})$ as a function of $\frac{\log(\frac{t}{\tau})}{\log(\frac{t_w}{\tau})}$ for AgMn at 8K. The values of $\tau = 10^{-5}$ sec. and $\frac{\theta}{\psi} = 0$ have been chosen to obtain the best rescaling.

Figure 8 $\frac{M(t_w + t, t_w)}{q_{EA}(T)}$ as a function of $\frac{\log(\frac{t}{\tau(T)})}{\log(\frac{t_w}{\tau(T)})}$ for CrIn at various temperatures. The values of $\tau(T)$ sec. and $q_{EA}(T)$ have been chosen to obtain the best rescaling of the late part of the curve, from which we extract $\frac{\lambda}{\psi} \simeq 1.2$.

Tables Extracted values of x_1 , x_2 , q_{EA} and q_1 , from CrMn and AgMn TRM decay data (using simple form of the FDT).

Appendix A - Some properties of exponential times

Here we shall review and give proofs of some of the properties of exponential times used in this paper. The probability density function of an exponential random variable T of rate E , is given by

$$p(t) = E \exp(-Et), \quad t \geq 0.$$

Property A-1: The exponential time is *memoryless*, that is conditional on the time T not having occurred at some time t , the distribution of the subsequent time T' before it occurs is the same as that for T . The proof uses Bayes' theorem to show that, the conditional probability density function for T' is

$$\rho(t') = \frac{E \exp(E(t + t'))}{\int_t^\infty ds E \exp(-Es)} = E \exp(-Et'),$$

and hence the result.

Property A-2: Given two independent exponential times T and T' of rates E and E' respectively, their minimum is distributed as an exponential random time of rate $E + E'$. This can be seen as follows: if the probability density function for $\min(T, T')$ is $\rho(u)$, then

$$\rho(u) = \int dt dt' E E' \exp(-Et - E't') \delta(u - \min(t, t')) = (E + E') \exp(-(E + E')u).$$

By induction we see that for any finite number of independent exponential times the minimum is distributed as an exponential with rate given by the sum of their rates.

Property A-3: Given two independent exponential times T and T' of rates E and E' respectively then

$$P(T < T') = \int_{t < t'} dt dt' EE' \exp(-Et - E't') = \frac{E}{E + E'}.$$

This property generalises to: for j independent exponential times T_i $1 \leq i \leq j$ of rates E_i $1 \leq i \leq j$ respectively, then

$$P(T_k = \min\{T_i, \quad 1 \leq i \leq j\}) = \frac{E_k}{\sum_{i=1}^j E_i}.$$

The proof simply uses the basic form of property (A-3) and also property (A-2).

REFERENCES

- [1] L.C.E. Struick, “Physical Aging in Amorphous Polymers and Other Materials” (Elsevier, Houston, 1978)
- [2] L. Lundgren, P. Svedlindh, P. Nordblad, O. Beckmann, Phys. Rev. Lett **51**, 911 (1983),
P. Nordblad, L. Lundgren, P. Svedlindh, L. Sandlund, Phys. Rev. **B 33**, 645 (1988)
- [3] M. Alba, M. Ocio, J. Hammann, Europhys. Lett. **2**, 45 (1986), J. Phys. Lett. **46** L-1101 (1985),
M. Alba, J. Hammann, M. Ocio, Ph. Refregier, J. Appl. Phys. 61, 3683 (1987),
E. Vincent, J. Hammann, M. Ocio, p. 207 in ”Recent Progress in Random Magnets”,
D.H. Ryan Editor, (World Scientific Pub. Co. Pte. Ltd, Singapore 1992)
- [4] V.S. Dotsenko, M. V. Feigelmann, L.B. Ioffe, Spin-Glasses and related problems, Soviet Scientific Reviews, vol. 15 (Harwood, 1990)
- [5] L. Cugliandolo, J. Kurchan, Phys. Rev. Lett. **71** (1993) 173
- [6] S. Franz, M. Mézard, Europhys. Lett. **26** (1994) 209, Physica **A209** (1994) 1
- [7] L. Cugliandolo, J. Kurchan, *On the out of Equilibrium Relaxation of the Sherrington Kirkpatrick model*, cond-mat/9311016 (1993)
- [8] L. Cugliandolo, J. Kurchan, G. Parisi, *Off equilibrium dynamics and aging in unfrustrated systems*, cond-mat/9406053 (1993)
- [9] J.O. Andersson, J. Mattson, P. Svedlindh, Phys. Rev. B **46** 8297 (1992)
- [10] H. Rieger, J. Phys. A **26** L615 (1993), H. Rieger, B. Steckemetz and M. Schreckenberg, Europhysics Letters, in press (1994).
- [11] L. Cugliandolo, J. Kurchan, F. Ritort, Phys. Rev. B **49**, 6331 (1994)
- [12] M. Mézard, G. Parisi, M.A. Virasoro, “Spin Glass Theory and Beyond”, (World Scientific, Singapore 1987)

- [13] B. Derrida, Phys. Rev. **B 24**, 2613 (1981)
- [14] a- J.P. Bouchaud, J. Physique I (France) **2**, 1705 (1992), b- J.P. Bouchaud, E. Vincent, J. Hammann, J. Physique I (France) **4**, 139 (1994)
- [15] D.J. Gross, M. Mézard, Nucl. Phys. **B240** (1984) 431.
- [16] There are quite a few recent papers on that subject: see e.g. J.P. Bouchaud, M. Mézard J. Phys. I France **4** (1994) 1109, E. Marinari, G.Parisi, F.Ritort, *Replica Field Theory for Deterministic Models: Binary Sequences with Low Autocorrelation I and II* preprint hep-th/9405148, ROM2F/94/15; preprint cond-mat/9406074, L. F. Cugliandolo, J. Kurchan, G.Parisi, F.Ritort, *Matrix Models as solvable Glass Models* preprint cond-mat 9407086 , S. Franz, J. Hertz, *Glassy transition and aging in a model without disorder* preprint cond-mat/9408075.
- [17] W. Krauth, M. Mézard, *Aging without disorder on long time scales*, LPENS preprint 94/20.
- [18] J. Hammann, M. Lederman, M. Ocio, R. Orbach, E. Vincent, Physica **A 185**, 278 (1992), F. Lefloch, J. Hammann, M. Ocio, E. Vincent, Europhys. Lett **18**, 647 (1992)
- [19] M. Schreckenberg, Z. Phys. B **60**, 483 (1985), C. Bachas, B. Huberman, Phys. Rev. Lett. **57** 1985 (1986); G. Paladin, M. Mézard, C. de Dominicis, J. Physique (Lettres) **46** L-985 (1985), R. G. Palmer, in ‘Chance and Matter’, Les Houches 1986, R. Stora, G, Toulouse, J. Vannimenus Edts, North Holland; P. Sibani, K.H. Hoffmann, Europhys. Lett. **4** 967 (1987) and refs. therein.
- [20] G. B. Alers, M. B. Weissmann, N.E. Isrealoff, Phys. Rev. **B 46**, 507 (1992), M. B. Weissmann, N.E. Isrealoff, G. B. Alers, Journal of Magn. Magn. Mat. **114**, 87 (1992), and M. B. Weissmann, Rev. Mod. Phys., July 1993.
- [21] D.S Fisher, D.A. Huse , Phys. Rev. Lett **56**, 1601 (1986), Phys. Rev. **B 38**, 373 (1988)

- [22] $x > 1$ indeed appears within Parisi's ansatz for some models, see e.g. J. Kurchan, G. Parisi, M.A. Virasoro, J. Phys. I (France) **3** (1993) 1819. The corresponding states are not sufficiently numerous to contribute to the Boltzmann weight, but certainly have a dynamical meaning.
- [23] The equilibrium dynamics (of the SK model) is described in H. Sompolinsky, A. Zippelius, Phys. Rev. B **25** 6960 (1982).
- [24] Note that for a ferromagnet with $H \neq 0$, both (11-a) and (11-b) are non zero.
- [25] P. Doussineau, Y. Farssi, C. Frénois, A. Levelut, J. Toulouse, S. Ziolkiewicz, J. Phys. I (France) **4** 1217 (1994). The dielectric response of $K_{1-y}LI_yTaO_3$ in the high temperature phase are well described by an [x]-distribution of relaxation time with $x \simeq \frac{T}{T_g} > 1$.
- [26] G. Parisi, in 'Chance and Matter', Les Houches 1986, R. Stora, G. Toulouse, J. Vannimenus Edts, North Holland.
- [27] see the discussion of the effective microscopic time appearing in E. Vincent, J. P. Bouchaud, D. S. Dean, J. Hammann, *Aging in spin glasses as a random walk: How does a magnetic field alter the landscape*, preprint.
- [28] We are indebted to Silvio Franz for an important discussion on this point.
- [29] The late time power law behaviour t^{-x_1} can be probed on nearly four decades using the attenuation of aging induced by a strong magnetic field : see [27]
- [30] K. Binder, A.P. Young, Rev. Mod. Phys. **58**, 801 (1986)
- [31] P. Refrégier, M. Ocio, Rev. Phys. Appl. **22** 367 (1987), P. Refrégier, Thèse, Université de Paris-Sud (1987).
- [32] G.J. Koper, H.J. Hilhorst, Physica A **155** 431 (1989)
- [33] J.O. Andersson, J. Mattson, P. Nordblad, Phys. Rev. B **48** 13977 (1993)

- [34] J.P. Bouchaud, D.S. Dean, J. Hammann, E. Vincent, in preparation.
- [35] F. Lefloch, Thèse N 2808, Université de Paris-Sud (1993)
- [36] We do not find the explanation based on the ‘droplet’ model (supplemented by chaoticity and ‘breaking time’ assumptions) [38], [33] fully satisfactory.
- [37] B. Derrida, J. Physique Lettres **46** 401 (1985), B. Derrida, E. Gardner, J. Phys. C **19** 2253 (1986).
- [38] G.J. Koper, H.J. Hilhorst, J. Physique (France) **49** 429 (1988)
- [39] A. J. Bray, *Theory of phase ordering kinetics*, to appear in Adv. Physics.
- [40] D.S. Dean, J.P. Bouchaud, in preparation.
- [41] J.P. Bouchaud, A. Georges, Phys. Rep. **195** 127 (1990), Chapter III
- [42] G. Parisi and E. Marinari, J. Phys. A **26**, L1149 (1993)
- [43] In a latter paper [32] the authors study analytically the 1d $\pm J$ spin glass. They indeed find $p \equiv \frac{1}{z} = \frac{1}{2}$, and compute $M(t_w + t, t_w)$. Interestingly, their result can be written *exactly* as in Eq. (9) with $x = \frac{1}{2}$. This is not a coincidence: the probability for a domain to survive during a time τ is given by the probability for two random walkers (the domain walls) to meet for the first time after time τ , which is, in 1d and for large τ , $\propto \tau^{-1-\frac{1}{2}}$.
- [44] M. Mézard, G. Parisi, J. Physique I **1** 809 (1991)
- [45] J.P. Bouchaud, M. Mézard, J. Yedidia, Phys. Rev B **46** 14 686 (1992)
- [46] T. Odagaki, J. Matsui, Y. Hiwatari, Physica A **204** 464 (1994).